# Gas in debris disks A new way to produce patterns?

# Wladimir Lyra

Sagan Fellow NASA Jet Propulsion Laboratory





## Marc Kuchner

NASA Goddard Space Flight Center

## <u>Debris disks</u>



## Sharp and eccentric rings in debris disks: Signposts of planets



## Narrow sharp eccentric ring

Detection of a source quickly heralded as a planet Fomalhaut b

## Sharp and eccentric rings in debris disks: Signposts of planets ?

## However....

THE ASTROPHYSICAL JOURNAL, 747:116 (7pp), 2012 March 10 © 2012. The American Autonomical Society. All rights merved. Printed in the U.S.A. dol:10.1088/0004-637X/747/2/116

# Fomalhaut De

Ring Oute NASA, ESA, P. Kala and M. Clampin (NA

#### INFRARED NON-DETECTION OF FOMALHAUT b: IMPLICATIONS FOR THE PLANET INTERPRETATION

MARKUS JANSON<sup>1,5</sup>, JOSEPH C. CARSON<sup>2</sup>, DAVID LAFRENIÈRE<sup>3</sup>, DAVID S. SPIEGEL<sup>4</sup>, JOHN R. BENT<sup>2</sup>, AND PALMER WONG<sup>2</sup> <sup>1</sup>Department of Astrophysical Sciences, Princeton University, Princeton, NJ, USA; Janson@astro.princeton.edu <sup>2</sup>College of Charleston, Charleston, WV, USA <sup>3</sup>Department of Physics, University of Montreal, Montreal, Canada <sup>4</sup>Institute for Advanced Studies, Princeton, NJ, USA *Received 2011 December 16*; accepted 2012 January 12; published 2012 February 23

#### ABSTRACT

The nearby A4-type star Fornalhaut hosts a debris belt in the form of an eccentric ring, which is thought to be caused by dynamical influence from a giant planet companion. In 2008, a detection of a point source inside the inner edge of the ring was reported and was interpreted as a direct image of the planet, named Fornalhaut b. The detection was made at ~600–800 nm, but no corresponding signatures were found in the near-infrared range, where the bulk emission of such a planet should be expected. Here, we present deep observations of Fornalhaut with *Spitzer*/IRAC at 4.5  $\mu$ m, using a novel point-spread function subtraction technicue based on angular differential imaging and Locally Optimized Combination of Images, in order to substantially improve the *Spitzer* contrast at small separations. The results provide more than an order of magnitude improvement in the upper flux limit of Fornalhaut b and exclude the possibility that any flux from a giant planet surface contributes to the observed flux at visible wavelengths. This renders any direct connection between the observed light source and the dynamically inferred giant planet highly unlikely. We discuss several possible interpretations of the total body of observations of the Fornalhaut system and find that the interpretation that best matches the available data for the observed source is scattered light from a transient or semi-transient dust cloud.

Key words: circumstellar matter - planetary systems - stars: early-type

Online-only material: color figures

## Planet not detected in infrared



## Sharp and eccentric rings in debris disks: Signposts of planets ?

## However.....

Fomalhaut Deb

No Data

Ring Oute

NASA, ESA, P. Kalas

and M. Clampin (NAS

24 Oct 2012

h.EP1



Thayne Currie<sup>1,2</sup>, John Debes<sup>3</sup>, Timothy J. Rodigas<sup>4</sup>, Adam Burrows<sup>5</sup>, Yoichi Itoh<sup>6</sup>, Misato Fukagawa<sup>7</sup>, Scott J. Kenyon<sup>8</sup>, Marc Kuchner<sup>2</sup>, Soko Matsumura<sup>9</sup>

currie@astro.utoronto.ca

t b Planet

STSH-PRC08-39

#### ABSTRACT

We present Subaru/IRCS J band data for Fomalhaut and a (re)reduction of archival 2004–2006 HST/ACS data first presented by Kalas et al. (2008). We confirm the existence of a candidate exoplanet, Fomalhaut b, in both the 2004 and 2006 F606W data sets at a high signal-to-noise. Additionally, we confirm



## Some of the Fom b controversy

## Janson et al. 2011 Boley et al. 2012

Variability by 0.7-0.8 mag in F606W band

Astrometric orbit not apsidally aligned with the ring

No infrared emission

Not seen by ALMA

Emission in optical required scattering by cloud of several Jupiter radii

Outer ring edge also sharp

### Currie et al. 2012

No variability found within 0.15 mag in the same band

Consistent with apsidal alignment

Thermal emission from a 0.5  $M_J$  planet (enough to sculpt the debris disk) would not be detectable anyway.

# Are there alternative explanations?

## Debris disks are not completely gas-free



## Gas in debris disks

<b>D21</b>	<b>ISCT</b>	IONS
$\overline{}$		

β Pictoris	many species	Lagrange et al. (1998),
51 Ophiuchi	many species	Roberge et al. (2002)
σHerculis	CII, NII	Chen & Jura (2003)
HD 32297	Na I, CII	Redfield (2007), Donaldson et al. (2012)
HD 135344	H2, CO	Thi et al. (2001), Pontoppidan et al. (2008)
49 Ceti	H2, CO	Dent et al. (2005), Roberge et al. (2012)
AU Mic H2	H2	France et al. (2007)
HD172555	510	Lisse et al. (2009)





Klahr & Lin (2005)

Suggested that an instability might cause dust in debris disks to clump together. Dust pressure trap



Adapted from Whipple (1972)

Dust pressure trap



## Dust pressure trap



Photoelectric heating

In optically thin debris disks, the **dust** is the **main heating agent** for the gas.



Dust intercepts starlight directly, emits electron, that heats the gas.

Gas is photoelectrically heated by the dust itself

Runaway process: instability

Dust heats gas Heated gas = high pressure region High pressure concentrates dust Runaway process: instability



## Dust heats gas

Heated gas = high pressure region High pressure concentrates dust



## Model equations

Klahr & Lin (2005) used a simplified, 1-D model.

## Model equations

Our simulation adds much more physics, and works in 2D.

$$\begin{split} & \mathsf{Klahr} \& \mathsf{Lin} (2005) \\ & \mathsf{1D} \end{split} \\ & \frac{\partial}{\partial t} \Sigma_d + \frac{1}{r} \frac{\partial}{\partial r} r \Sigma_d v_r = 0. \\ & V_\phi = \Omega r + \frac{1}{2\Omega \Sigma_g} \frac{\partial}{\partial r} P \\ & T_g = T_0 \left( \frac{\Sigma_d}{\Sigma_0} \right)^\beta, \end{split}$$

Inertia for both gas and dust

Energy equation

Drag force and drag force backreaction

Lyra & Kuchner (2012)  

$$\frac{\partial \Sigma_g}{\partial t} = -(u \cdot \nabla) \Sigma_g - \Sigma_g \nabla \cdot u$$

$$\frac{\partial u}{\partial t} = -(u \cdot \nabla) u - \frac{1}{\Sigma_g} \nabla P - \nabla \Phi - \frac{\Sigma_d}{\Sigma_g} f_d$$

$$\frac{\partial S}{\partial t} = -(u \cdot \nabla) S - \frac{c_v}{T} \frac{(T - T_p)}{\tau_T}.$$

$$\frac{dx}{dt} = v$$

$$\frac{dv}{dt} = -\nabla \Phi + f_d$$

$$f_d = -\frac{(v - u)}{\tau_f}$$

$$T_p = T_0 \frac{\Sigma_d}{\Sigma_0}.$$

## Linear Analysis







## Linear and nonlinear growth



Linear growth only exists for  $\epsilon < 1$ 

But there is nonlinear growth beyond !

## Instability



Narrow hot dust rings Cold gas collects between rings Ring width

Ring spacing and width is determined by the wavelength of maximum growth.



Which in turn is determined by viscosity

Ring width ~ 10 Kolmogorov lengths

## Robustness



## Oscillations





## Solutions



## Solutions



Damped and free Oscillations

## Solutions



#### Damped and free Oscillations



## The model in 2D: Eccentric rings



Azimuth

Epicyclic oscillations

make the ring appear *eccentric* !!!



## Ring eccentricity



## Thermocentrifugal instability in Debris Disks with Gas:

## Driven by pressure trapping + photoelectric heating.



## Thermocentrifugal instability in Debris Disks with Gas:

Driven by pressure trapping + photoelectric heating.

## No radial drift, and axissymetric linear modes. Nothing to do with the streaming instability

Robust to inertia, drag backreaction, thermal time, friction time

Rings of width ~10  $\lambda_{\mu}$  ( $\lambda_{\mu}$  ~ 1 AU for Fomalhaut at 100 AU)

Linear for dust-to-gas ratio <= 1, nonlinear beyond

Produces sharp narrow eccentric rings. Caution before shouting "Planet!". Not all that glitters is gold...

Future Work:

3D Kinetic theory Interplay with streaming instability Extra physics: Photophoresis, Collisions, Magnetic Fields

## Thermocentrifugal instability in Debris Disks with Gas:

Driven by pressure trapping + photoelectric heating.

No radial drift, and axissymetric linear modes. Nothing to do with the streaming instability

## Robust to inertia, drag backreaction, thermal time, friction time



## Thermocentrifugal instability in Debris Disks with Gas:

Driven by pressure trapping + photoelectric heating.

No radial drift, and axissymetric linear modes. Nothing to do with the streaming instability

Robust to inertia, drag backreaction, thermal time, friction time

## Rings of width ~10 $\lambda_{v}$ ( $\lambda_{v}$ ~ 1 AU for Fomalhaut at 100 AU)



## Thermocentrifugal instability in Debris Disks with Gas:

Driven by pressure trapping + photoelectric heating.

No radial drift, and axissymetric linear modes. Nothing to do with the streaming instability

Robust to inertia, drag backreaction, thermal time, friction time

Rings of width ~10  $\lambda_{\mu}$  ( $\lambda_{\mu}$  ~ 1 AU for Fomalhaut at 100 AU)

## Linear for dust-to-gas ratio <= 1, nonlinear beyond



## Thermocentrifugal instability in Debris Disks with Gas:



Produces sharp narrow eccentric rings. Caution before shouting "Planet!". Not all that glitters is gold...

**Future Work:** 

3D Kinetic theory Interplay with streaming instability Extra physics: Photophoresis, Collisions, Magnetic Fields

Thermocentrifugal instability in Debris Disks with Gas:

Driven by pressure trapping + photoelectric heating.

No radial drift, and axissymetric linear modes. Nothing to do with the streaming instability

Robust to inertia, drag backreaction, thermal time, friction time

Rings of width ~10  $\lambda_{\mu}$  ( $\lambda_{\mu}$  ~ 1 AU for Fomalhaut at 100 AU)

Linear for dust-to-gas ratio <= 1, nonlinear beyond

Produces sharp narrow eccentric rings. Caution before shouting "Planet!". Not all that glitters is gold...

## **Future Work:**

3D Kinetic theory Interplay with streaming instability Extra physics: Photophoresis, Collisions, Magnetic Fields

## arXiv:1204.6322

DRAFT VERSION Preprint typeset using LATEX style emulateapj v. 12/16/11

#### SHARP ECCENTRIC RINGS IN PLANETLESS HYDRODYNAMICAL MODELS OF DEBRIS DISKS.

WLADIMIR LYRA<sup>1,2,3</sup> AND MARC J. KUCHNER<sup>4</sup>

Draft version

#### ABSTRACT

Debris disks should not be completely gas-free, since there is second generation gas from outgassing of planetesimals and dust grains via sublimation, photodesorption, or collisions, generating a system of dust-to-gas ratio close to unity, where hydrodynamics cannot be ignored. A clumping instability exists in this configuration, that has been hitherto explored only in one-dimensional, incompressible models. We performed 2D numerical compressible models of a disk with comparable amounts of gas and dust to study the growth and development of this instability. Our model solves the momentum equation for the gas and dust, together with energy and continuity equations. We uncover that the backreaction of the drag force from the gas onto the dust shepherds rings, similar to those observed in debris disks and usually attributed to the presence of hypothetical undetected planets. We also uncover that the eccentricity of these rings, usually presented as convincing evidence for the presence of a planet, can actually be simply explained by a standing wave propagating along the ring. The rings support a spectrum of oscillations, with one particular mode representing epicyclic motion. The apparent eccentricity matches the eccentricity in observed systems. This suggests that the planet possibility, though thrilling, is not necessarily required to explain these systems.

#### 1. INTRODUCTION

Disks around young stars appear to pass through an evolutionary phase when the disk is optically-thin and the dust to gas ratio is of order unity give or take an order of magnitude. It can be hard to precisely measure the total masses of the dust and gas in such disks, but the nearby stars  $\beta$  Pictoris (Lagrange et al. 1998; Olofsson et al. 2001; Brandeker et al. 2004; Roberge et al. 2006; Troutman et al. 2011), HD32297 (Redfield 2007), 49 Ceti (Zuckerman et al. 1995) and HD 21997 (Moor et al. 2011) all host disks of dust resembling ordinary debris disks and also have stable circumstellar gas detected in molecular CO, Na I or other metal lines; the inferred mass of

The result of this instability could be that the dust and gas clump into rings or spiral patterns or other structures that could be detected via coronographic imaging or other methods. Indeed, images of debris disks and transitional disks show a range of asymmetries and other structures that beg for explanation. Klahr & Lin (2005) raised the possibility that the instability they hypothesized could explain some of the observed structures. Alternative explanations for these structures sometimes rely on planetary perturbers–a tantalizing possibility. But we are interested in investigating any possible explanation for these disk structures that does not require a hidden planetary companion.

## arXiv:1204.6322



## SHARP ECCENTRIC RINGS IN PLANETLESS HYDRODYNAMICAL MODELS OF DEBRIS DISKS.





(Zuckerman et al. 1995) and HD 21997 (Moor et al. 2011) all host disks of dust resembling ordinary debris disks and also have stable circumstellar gas detected in molecular CO, Na I or other metal lines; the inferred mass of sometimes rely on planetary perturbers-a tantalizing possibility. But we are interested in investigating any possible explanation for these disk structures that does not require a hidden planetary companion.

 $M_{\rm eff}$  (1997)  $M_{\rm eff}$ 

